

# Package ‘correlation’

September 27, 2020

**Type** Package

**Title** Methods for Correlation Analysis

**Version** 0.4.0

**Maintainer** Dominique Makowski <dom.makowski@gmail.com>

**URL** <https://easystats.github.io/correlation/>

**BugReports** <https://github.com/easystats/correlation/issues>

**Description** Lightweight package for computing different kinds of correlations, such as partial correlations, Bayesian correlations, multilevel correlations, polychoric correlations, biweight correlations, distance correlations and more. Relies on the easystats ecosystem (Lüdecke, Waggoner & Makowski (2019) <doi:10.21105/joss.01412>).

**Imports** datasets, stats, insight (>= 0.8.0), bayestestR (>= 0.5.0), parameters (>= 0.5.0), effectsize (>= 0.2.0)

**Suggests** BayesFactor, dplyr, energy, forcats, ggcorrplot, ggplot2, Hmisc, knitr, lme4, polycor, ppcor, psych, rmcrr, testthat, tidy, covr, rstanarm, rmarkdown, see, wdm, WRS2

**License** GPL-3

**Encoding** UTF-8

**LazyData** true

**RoxygenNote** 7.1.1

**VignetteBuilder** knitr

**NeedsCompilation** no

**Author** Dominique Makowski [aut, cre] (<<https://orcid.org/0000-0001-5375-9967>>),  
Daniel Lüdecke [aut] (<<https://orcid.org/0000-0002-8895-3206>>),  
Mattan S. Ben-Shachar [ctb] (<<https://orcid.org/0000-0002-4287-4801>>),  
Indrajeet Patil [ctb] (<<https://orcid.org/0000-0003-1995-6531>>)

**Repository** CRAN

**Date/Publication** 2020-09-27 10:40:02 UTC

**R topics documented:**

correlation . . . . .	2
cor_test . . . . .	6
cor_to_ci . . . . .	10
cor_to_cov . . . . .	11
cor_to_pcor . . . . .	12
distance_mahalanobis . . . . .	13
is.cor . . . . .	14
isSquare . . . . .	14
matrix_inverse . . . . .	15
simulate_simpson . . . . .	15
z_fisher . . . . .	16
<b>Index</b>	<b>18</b>

---

correlation	<i>Correlation Analysis</i>
-------------	-----------------------------

---

**Description**

Performs a correlation analysis.

**Usage**

```
correlation(
  data,
  data2 = NULL,
  method = "pearson",
  p_adjust = "holm",
  ci = 0.95,
  bayesian = FALSE,
  bayesian_prior = "medium",
  bayesian_ci_method = "hdi",
  bayesian_test = c("pd", "rope", "bf"),
  redundant = FALSE,
  include_factors = FALSE,
  partial = FALSE,
  partial_bayesian = FALSE,
  multilevel = FALSE,
  robust = FALSE,
  ...
)
```

**Arguments**

data	A data frame.
data2	An optional data frame.
method	A character string indicating which correlation coefficient is to be used for the test. One of "pearson" (default), "kendall", or "spearman", "biserial", "polychoric", "tetrachoric", "biweight", "distance", "percentage" (for percentage bend correlation), "blomqvist" (for Blomqvist's coefficient), "hoeffding" (for Hoeffding's D), "gamma", "gaussian" (for Gaussian Rank correlation) or "shepherd" (for Shepherd's Pi correlation). Setting "auto" will attempt at selecting the most relevant method (polychoric when ordinal factors involved, tetrachoric when dichotomous factors involved, point-biserial if one dichotomous and one continuous and pearson otherwise).
p_adjust	Correction method for frequentist correlations. Can be one of "holm" (default), "hochberg", "hommel", "bonferroni", "BH", "BY", "fdr" or "none".
ci	Confidence/Credible Interval level. If "default", then it is set to 0.95 (95% CI).
bayesian	If TRUE, will run the correlations under a Bayesian framework. Note that for partial correlations, you will also need to set <code>partial_bayesian</code> to TRUE to obtain "full" Bayesian partial correlations. Otherwise, you will obtain pseudo-Bayesian partial correlations (i.e., Bayesian correlation based on frequentist partialization).
bayesian_prior	For the prior argument, several named values are recognized: "medium.narrow", "medium", "wide", and "ultrawide". These correspond to scale values of $1/\sqrt{27}$ , $1/3$ , $1/\sqrt{3}$ and 1, respectively. See the <code>BayesFactor::correlationBF</code> function.
bayesian_ci_method	See arguments in <a href="#">model_parameters</a> for BayesFactor tests.
bayesian_test	See arguments in <a href="#">model_parameters</a> for BayesFactor tests.
redundant	Should the data include redundant rows (where each given correlation is repeated two times).
include_factors	If TRUE, the factors are kept and eventually converted to numeric or used as random effects (depending of <code>multilevel</code> ). If FALSE, factors are removed upfront.
partial	Can be TRUE or "semi" for partial and semi-partial correlations, respectively.
partial_bayesian	If TRUE, will run the correlations under a Bayesian framework. Note that for partial correlations, you will also need to set <code>partial_bayesian</code> to TRUE to obtain "full" Bayesian partial correlations. Otherwise, you will obtain pseudo-Bayesian partial correlations (i.e., Bayesian correlation based on frequentist partialization).
multilevel	If TRUE, the factors are included as random factors. Else, if FALSE (default), they are included as fixed effects in the simple regression model.
robust	If TRUE, will rank-transform the variables prior to estimating the correlation. Note that, for instance, a Pearson's correlation on rank-transformed data is equivalent to a Spearman's rank correlation. Thus, using <code>robust=TRUE</code> and <code>method="spearman"</code>

is redundant. Nonetheless, it is an easy way to increase the robustness of the correlation (as well as obtaining Bayesian Spearman rank Correlations).

... Arguments passed to or from other methods.

## Details

### Correlation Types:

- **Pearson's correlation:** This is the most common correlation method. It corresponds to the covariance of the two variables normalized (i.e., divided) by the product of their standard deviations.
- **Spearman's rank correlation:** A non-parametric measure of rank correlation (statistical dependence between the rankings of two variables). The Spearman correlation between two variables is equal to the Pearson correlation between the rank values of those two variables; while Pearson's correlation assesses linear relationships, Spearman's correlation assesses monotonic relationships (whether linear or not). Confidence Intervals (CI) for Spearman's correlations are computed using the Fieller et al. (1957) correction (see Bishara and Hittner, 2017).
- **Kendall's rank correlation:** In the normal case, the Kendall correlation is preferred than the Spearman correlation because of a smaller gross error sensitivity (GES) and a smaller asymptotic variance (AV), making it more robust and more efficient. However, the interpretation of Kendall's tau is less direct than that of Spearman's rho, in the sense that it quantifies the difference between the % of concordant and discordant pairs among all possible pairwise events. Confidence Intervals (CI) for Kendall's correlations are computed using the Fieller et al. (1957) correction (see Bishara and Hittner, 2017).
- **Biweight midcorrelation:** A measure of similarity that is median-based, instead of the traditional mean-based, thus being less sensitive to outliers. It can be used as a robust alternative to other similarity metrics, such as Pearson correlation (Langfelder & Horvath, 2012).
- **Distance correlation:** Distance correlation measures both linear and non-linear association between two random variables or random vectors. This is in contrast to Pearson's correlation, which can only detect linear association between two random variables.
- **Percentage bend correlation:** Introduced by Wilcox (1994), it is based on a down-weight of a specified percentage of marginal observations deviating from the median (by default, 20%).
- **Shepherd's Pi correlation:** Equivalent to a Spearman's rank correlation after outliers removal (by means of bootstrapped Mahalanobis distance).
- **Blomqvist's coefficient:** The Blomqvist's coefficient (also referred to as Blomqvist's Beta or medial correlation; Blomqvist, 1950) is a median-based non-parametric correlation that has some advantages over measures such as Spearman's or Kendall's estimates (see Schmid and Schimdt, 2006).
- **Hoeffding's D:** The Hoeffding's D statistic is a non-parametric rank based measure of association that detects more general departures from independence (Hoeffding 1948), including non-linear associations. Hoeffding's D varies between -0.5 and 1 (if there are no tied ranks, otherwise it can have lower values), with larger values indicating a stronger relationship between the variables.
- **Point-Biserial and biserial correlation:** Correlation coefficient used when one variable is continuous and the other is dichotomous (binary). Point-Biserial is equivalent to a Pearson's correlation, while Biserial should be used when the binary variable is assumed to have an

underlying continuity. For example, anxiety level can be measured on a continuous scale, but can be classified dichotomously as high/low.

- **Gamma correlation:** The Goodman-Kruskal gamma statistic is similar to Kendall's Tau coefficient. It is relatively robust to outliers and deals well with data that have many ties.
- **Gaussian rank Correlation:** The Gaussian rank correlation estimator is a simple and well-performing alternative for robust rank correlations (Boudt et al., 2012). It is based on the Gaussian quantiles of the ranks.
- **Polychoric correlation:** Correlation between two theorised normally distributed continuous latent variables, from two observed ordinal variables.
- **Tetrachoric correlation:** Special case of the polychoric correlation applicable when both observed variables are dichotomous.

**Partial Correlation:** **Partial correlations** are estimated as the correlation between two variables after adjusting for the (linear) effect of one or more other variable. The correlation test is then run after having partialized the dataset, independently from it. In other words, it considers partialization as an independent step generating a different dataset, rather than belonging to the same model. This is why some discrepancies are to be expected for the t- and p-values, CIs, BFs etc (but *not* the correlation coefficient) compared to other implementations (e.g., `ppcor`). (The size of these discrepancies depends on the number of covariates partialled-out and the strength of the linear association between all variables.) Such partial correlations can be represented as Gaussian Graphical Models (GGM), an increasingly popular tool in psychology. A GGM traditionally include a set of variables depicted as circles ("nodes"), and a set of lines that visualize relationships between them, which thickness represents the strength of association (see Bhushan et al., 2019).

**Multilevel correlations** are a special case of partial correlations where the variable to be adjusted for is a factor and is included as a random effect in a mixed model.

#### Notes:

- Kendall and Spearman correlations when `bayesian=TRUE`: These are technically Pearson Bayesian correlations of rank transformed data, rather than pure Bayesian rank correlations (which have different priors).

#### Value

A correlation object that can be displayed using the `print`, `summary` or `table` methods.

**Multiple tests correction:** About multiple tests corrections.

#### References

- Boudt, K., Cornelissen, J., & Croux, C. (2012). The Gaussian rank correlation estimator: robustness properties. *Statistics and Computing*, 22(2), 471-483.
- Bhushan, N., Mohnert, F., Sloot, D., Jans, L., Albers, C., & Steg, L. (2019). Using a Gaussian graphical model to explore relationships between items and variables in environmental psychology research. *Frontiers in psychology*, 10, 1050.
- Bishara, A. J., & Hittner, J. B. (2017). Confidence intervals for correlations when data are not normal. *Behavior research methods*, 49(1), 294-309.

- Fieller, E. C., Hartley, H. O., & Pearson, E. S. (1957). Tests for rank correlation coefficients. I. *Biometrika*, 44(3/4), 470-481.
- Langfelder, P., & Horvath, S. (2012). Fast R functions for robust correlations and hierarchical clustering. *Journal of statistical software*, 46(11).
- Blomqvist, N. (1950). On a measure of dependence between two random variables, *Annals of Mathematical Statistics*, 21, 593-600

### Examples

```
library(correlation)
results <- correlation(iris)

results
summary(results)
summary(results, redundant = TRUE)

# Grouped dataframe
if (require("dplyr")) {
  iris %>%
    group_by(Species) %>%
    correlation()
}

# automatic selection of correlation method
correlation(mtcars[-2], method = "auto")
```

---

cor\_test

*Correlation test*

---

### Description

This function performs a correlation test between two variables.

### Usage

```
cor_test(
  data,
  x,
  y,
  method = "pearson",
  ci = 0.95,
  bayesian = FALSE,
  bayesian_prior = "medium",
  bayesian_ci_method = "hdi",
  bayesian_test = c("pd", "rope", "bf"),
  include_factors = FALSE,
  partial = FALSE,
  partial_bayesian = FALSE,
```

```

    multilevel = FALSE,
    robust = FALSE,
    ...
)

```

## Arguments

data	A data frame.
x, y	Names of two variables present in the data.
method	A character string indicating which correlation coefficient is to be used for the test. One of "pearson" (default), "kendall", or "spearman", "biserial", "polychoric", "tetrachoric", "biweight", "distance", "percentage" (for percentage bend correlation), "blomqvist" (for Blomqvist's coefficient), "hoeffding" (for Hoeffding's D), "gamma", "gaussian" (for Gaussian Rank correlation) or "shepherd" (for Shepherd's Pi correlation). Setting "auto" will attempt at selecting the most relevant method (polychoric when ordinal factors involved, tetrachoric when dichotomous factors involved, point-biserial if one dichotomous and one continuous and pearson otherwise).
ci	Confidence/Credible Interval level. If "default", then it is set to 0.95 (95% CI).
bayesian, partial_bayesian	If TRUE, will run the correlations under a Bayesian framework. Note that for partial correlations, you will also need to set partial_bayesian to TRUE to obtain "full" Bayesian partial correlations. Otherwise, you will obtain pseudo-Bayesian partial correlations (i.e., Bayesian correlation based on frequentist partialization).
bayesian_prior	For the prior argument, several named values are recognized: "medium.narrow", "medium", "wide", and "ultrawide". These correspond to scale values of $1/\sqrt{27}$ , $1/3$ , $1/\sqrt{3}$ and 1, respectively. See the <code>BayesFactor::correlationBF</code> function.
bayesian_ci_method, bayesian_test	See arguments in <a href="#">model_parameters</a> for BayesFactor tests.
include_factors	If TRUE, the factors are kept and eventually converted to numeric or used as random effects (depending of multilevel). If FALSE, factors are removed upfront.
partial	Can be TRUE or "semi" for partial and semi-partial correlations, respectively.
multilevel	If TRUE, the factors are included as random factors. Else, if FALSE (default), they are included as fixed effects in the simple regression model.
robust	If TRUE, will rank-transform the variables prior to estimating the correlation. Note that, for instance, a Pearson's correlation on rank-transformed data is equivalent to a Spearman's rank correlation. Thus, using robust=TRUE and method="spearman" is redundant. Nonetheless, it is an easy way to increase the robustness of the correlation (as well as obtaining Bayesian Spearman rank Correlations).
...	Arguments passed to or from other methods.

## Details

### Correlation Types:

- **Pearson's correlation:** This is the most common correlation method. It corresponds to the covariance of the two variables normalized (i.e., divided) by the product of their standard deviations.
- **Spearman's rank correlation:** A non-parametric measure of rank correlation (statistical dependence between the rankings of two variables). The Spearman correlation between two variables is equal to the Pearson correlation between the rank values of those two variables; while Pearson's correlation assesses linear relationships, Spearman's correlation assesses monotonic relationships (whether linear or not). Confidence Intervals (CI) for Spearman's correlations are computed using the Fieller et al. (1957) correction (see Bishara and Hittner, 2017).
- **Kendall's rank correlation:** In the normal case, the Kendall correlation is preferred than the Spearman correlation because of a smaller gross error sensitivity (GES) and a smaller asymptotic variance (AV), making it more robust and more efficient. However, the interpretation of Kendall's tau is less direct than that of Spearman's rho, in the sense that it quantifies the difference between the % of concordant and discordant pairs among all possible pairwise events. Confidence Intervals (CI) for Kendall's correlations are computed using the Fieller et al. (1957) correction (see Bishara and Hittner, 2017).
- **Biweight midcorrelation:** A measure of similarity that is median-based, instead of the traditional mean-based, thus being less sensitive to outliers. It can be used as a robust alternative to other similarity metrics, such as Pearson correlation (Langfelder & Horvath, 2012).
- **Distance correlation:** Distance correlation measures both linear and non-linear association between two random variables or random vectors. This is in contrast to Pearson's correlation, which can only detect linear association between two random variables.
- **Percentage bend correlation:** Introduced by Wilcox (1994), it is based on a down-weight of a specified percentage of marginal observations deviating from the median (by default, 20%).
- **Shepherd's Pi correlation:** Equivalent to a Spearman's rank correlation after outliers removal (by means of bootstrapped Mahalanobis distance).
- **Blomqvist's coefficient:** The Blomqvist's coefficient (also referred to as Blomqvist's Beta or medial correlation; Blomqvist, 1950) is a median-based non-parametric correlation that has some advantages over measures such as Spearman's or Kendall's estimates (see Schmid and Schimdt, 2006).
- **Hoeffding's D:** The Hoeffding's D statistic is a non-parametric rank based measure of association that detects more general departures from independence (Hoeffding 1948), including non-linear associations. Hoeffding's D varies between -0.5 and 1 (if there are no tied ranks, otherwise it can have lower values), with larger values indicating a stronger relationship between the variables.
- **Point-Biserial and biserial correlation:** Correlation coefficient used when one variable is continuous and the other is dichotomous (binary). Point-Biserial is equivalent to a Pearson's correlation, while Biserial should be used when the binary variable is assumed to have an underlying continuity. For example, anxiety level can be measured on a continuous scale, but can be classified dichotomously as high/low.
- **Gamma correlation:** The Goodman-Kruskal gamma statistic is similar to Kendall's Tau coefficient. It is relatively robust to outliers and deals well with data that have many ties.



- **Gaussian rank Correlation:** The Gaussian rank correlation estimator is a simple and well-performing alternative for robust rank correlations (Boudt et al., 2012). It is based on the Gaussian quantiles of the ranks.
- **Polychoric correlation:** Correlation between two theorised normally distributed continuous latent variables, from two observed ordinal variables.
- **Tetrachoric correlation:** Special case of the polychoric correlation applicable when both observed variables are dichotomous.

**Partial Correlation:** **Partial correlations** are estimated as the correlation between two variables after adjusting for the (linear) effect of one or more other variable. The correlation test is then run after having partialized the dataset, independently from it. In other words, it considers partialization as an independent step generating a different dataset, rather than belonging to the same model. This is why some discrepancies are to be expected for the t- and p-values, CIs, BFs etc (but *not* the correlation coefficient) compared to other implementations (e.g., ppcor). (The size of these discrepancies depends on the number of covariates partialled-out and the strength of the linear association between all variables.) Such partial correlations can be represented as Gaussian Graphical Models (GGM), an increasingly popular tool in psychology. A GGM traditionally include a set of variables depicted as circles ("nodes"), and a set of lines that visualize relationships between them, which thickness represents the strength of association (see Bhushan et al., 2019).

**Multilevel correlations** are a special case of partial correlations where the variable to be adjusted for is a factor and is included as a random effect in a mixed model.

#### Notes:

- Kendall and Spearman correlations when bayesian=TRUE: These are technically Pearson Bayesian correlations of rank transformed data, rather than pure Bayesian rank correlations (which have different priors).

#### Examples

```
library(correlation)

cor_test(iris, "Sepal.Length", "Sepal.Width")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "spearman")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "kendall")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "biweight")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "distance")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "percentage")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "blomqvist")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "hoeffding")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "gamma")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "gaussian")
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "shepherd")
cor_test(iris, "Sepal.Length", "Sepal.Width", bayesian = TRUE)

# Tetrachoric
data <- iris
data$Sepal.Width_binary <- ifelse(data$Sepal.Width > 3, 1, 0)
data$Petal.Width_binary <- ifelse(data$Petal.Width > 1.2, 1, 0)
cor_test(data, "Sepal.Width_binary", "Petal.Width_binary", method = "tetrachoric")
```

```

# Biserial
cor_test(data, "Sepal.Width", "Petal.Width_binary", method = "biserial")

# Polychoric
data$Petal.Width_ordinal <- as.factor(round(data$Petal.Width))
data$Sepal.Length_ordinal <- as.factor(round(data$Sepal.Length))
cor_test(data, "Petal.Width_ordinal", "Sepal.Length_ordinal", method = "polychoric")

# When one variable is continuous, will run 'polyserial' correlation
cor_test(data, "Sepal.Width", "Sepal.Length_ordinal", method = "polychoric")

# Robust (these two are equivalent)
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "pearson", robust = TRUE)
cor_test(iris, "Sepal.Length", "Sepal.Width", method = "spearman", robust = FALSE)

# Partial
cor_test(iris, "Sepal.Length", "Sepal.Width", partial = TRUE)
cor_test(iris, "Sepal.Length", "Sepal.Width", multilevel = TRUE)

cor_test(iris, "Sepal.Length", "Sepal.Width", partial_bayesian = TRUE)

```

---

cor\_to\_ci

---

*Convert correlation to p-values and CIs*


---

## Description

Get statistics, p-values and confidence intervals (CI) from correlation coefficients.

## Usage

```
cor_to_ci(cor, n, ci = 0.95, method = "pearson", correction = "fieller", ...)
```

```
cor_to_p(cor, n, method = "pearson")
```

## Arguments

cor	A correlation matrix or coefficient.
n	The sample size (number of observations).
ci	Confidence/Credible Interval level. If "default", then it is set to 0.95 (95% CI).
method	A character string indicating which correlation coefficient is to be used for the test. One of "pearson" (default), "kendall", or "spearman", "biserial", "polychoric", "tetrachoric", "biweight", "distance", "percentage" (for percentage bend correlation), "blomqvist" (for Blomqvist's coefficient), "hoeffding" (for Hoeffding's D), "gamma", "gaussian" (for Gaussian Rank correlation) or "shepherd" (for Shepherd's Pi correlation). Setting "auto" will attempt at selecting the most

	relevant method (polychoric when ordinal factors involved, tetrachoric when dichotomous factors involved, point-biserial if one dichotomous and one continuous and pearson otherwise).
correction	Only used if method is 'spearman' or 'kendall'. Can be 'fieller' (default; Fieller et al., 1957), 'bw' (only for Spearman) or 'none'. Bonett and Wright (2000) claim their correction ('bw') performs better, though the Bishara and Hittner (2017) paper favours the Fieller correction. Both are generally very similar.
...	Arguments passed to or from other methods.

**Value**

A list containing a p-value and the statistic or the CI bounds.

**References**

Bishara, A. J., & Hittner, J. B. (2017). Confidence intervals for correlations when data are not normal. *Behavior research methods*, 49(1), 294-309.

**Examples**

```
cor.test(iris$Sepal.Length, iris$Sepal.Width)
cor_to_p(-0.1175698, n = 150)
cor_to_p(cor(iris[1:4]), n = 150)
cor_to_ci(-0.1175698, n = 150)
cor_to_ci(cor(iris[1:4]), n = 150)

cor.test(iris$Sepal.Length, iris$Sepal.Width, method = "spearman")
cor_to_p(-0.1667777, n = 150, method = "spearman")
cor_to_ci(-0.1667777, ci = 0.95, n = 150)

cor.test(iris$Sepal.Length, iris$Sepal.Width, method = "kendall")
cor_to_p(-0.07699679, n = 150, method = "kendall")
```

---

cor\_to\_cov

---

*Convert a correlation to covariance*


---

**Description**

Convert a correlation to covariance

**Usage**

```
cor_to_cov(cor, sd = NULL, variance = NULL, tol = .Machine$double.eps^(2/3))
```

**Arguments**

cor	A correlation matrix, or a partial or a semipartial correlation matrix.
sd, variance	A vector that contains the standard deviations, or the variance, of the variables in the correlation matrix.
tol	Relative tolerance to detect zero singular values.

**Value**

A covariance matrix.

**Examples**

```
cor <- cor(iris[1:4])
cov(iris[1:4])

cor_to_cov(cor, sd = sapply(iris[1:4], sd))
cor_to_cov(cor, variance = sapply(iris[1:4], var))
```

---

 cor\_to\_pcor

---

*Correlation Matrix to (Semi) Partial Correlations*


---

**Description**

Convert a correlation matrix to a (semi)partial correlation matrix. Partial correlations are a measure of the correlation between two variables that remains after controlling for (i.e., "partialling" out) all the other relationships. They can be used for graphical Gaussian models, as they represent the direct interactions between two variables, conditioned on all remaining variables. This means that the squared partial correlation between a predictor X1 and a response variable Y can be interpreted as the proportion of (unique) variance accounted for by X1 relative to the residual or unexplained variance of Y that cannot be accounted for by the other variables.

**Usage**

```
cor_to_pcor(cor, tol = .Machine$double.eps^(2/3))

pcor_to_cor(pcor, tol = .Machine$double.eps^(2/3))

cor_to_spcor(cor = NULL, cov = NULL, tol = .Machine$double.eps^(2/3))

spcor_to_cor(
  spcor = NULL,
  cov = NULL,
  semi = FALSE,
  tol = .Machine$double.eps^(2/3)
)
```

**Arguments**

cor, pcor, spcor	A correlation matrix, or a partial or a semipartial correlation matrix.
tol	Relative tolerance to detect zero singular values.
cov	A covariance matrix (or a vector of the SD of the variables). Required for semi-partial correlations.
semi	Semi-partial correlations.

**Details**

The semi-partial correlation is similar to the partial correlation statistic. However, it represents (when squared) the proportion of (unique) variance accounted for by the predictor  $X_1$ , relative to the total variance of  $Y$ . Thus, it might be seen as a better indicator of the "practical relevance" of a predictor, because it is scaled to (i.e., relative to) the total variability in the response variable.

**Value**

The (semi) partial correlation matrix.

**Examples**

```
cor <- cor(iris[1:4])

# Partialize
cor_to_pcor(cor)
cor_to_spcor(cor, cov = sapply(iris[1:4], sd))

# Inverse
round(pcor_to_cor(cor_to_pcor(cor)) - cor, 2) # Should be 0
```

---

distance\_mahalanobis *Mahalanobis distance and confidence interval (CI)*

---

**Description**

The Mahalanobis distance (in squared units) measures the distance in multivariate space taking into account the covariance structure of the data. Because a few extreme outliers can skew the covariance estimate, the bootstrapped version is considered as more robust.

**Usage**

```
distance_mahalanobis(data, ci = 0.95, iterations = 1000, robust = TRUE, ...)
```

**Arguments**

data	A data frame.
ci	Confidence/Credible Interval level. If "default", then it is set to 0.95 (95% CI).
iterations	The number of draws to simulate/bootstrap (when robust is TRUE).
robust	If TRUE, will run a bootstrapped version of the function with $i$ iterations.
...	Arguments passed to or from other methods.

**Value**

Description of the Mahalanobis distance.

**References**

- Schwarzkopf, D. S., De Haas, B., & Rees, G. (2012). Better ways to improve standards in brain-behavior correlation analysis. *Frontiers in human neuroscience*, 6, 200.

**Examples**

```
distance_mahalanobis(iris[, 1:4])
distance_mahalanobis(iris[, 1:4], robust = FALSE)
```

---

is.cor	<i>Check if matrix resembles a correlation matrix</i>
--------	---

---

**Description**

Check if matrix resembles a correlation matrix

**Usage**

```
is.cor(x)
```

**Arguments**

x                    A matrix.

**Value**

TRUE of the matrix is a correlation matrix or FALSE otherwise.

---

isSquare	<i>Check if Square Matrix</i>
----------	-------------------------------

---

**Description**

Check if Square Matrix

**Usage**

```
isSquare(m)
```

**Arguments**

m                    A matrix.

**Value**

TRUE of the matrix is square or FALSE otherwise.

---

matrix_inverse	<i>Matrix Inversion</i>
----------------	-------------------------

---

**Description**

Performs a Moore-Penrose generalized inverse (also called the Pseudoinverse).

**Usage**

```
matrix_inverse(m, tol = .Machine$double.eps^(2/3))
```

**Arguments**

m	Matrix for which the inverse is required.
tol	Relative tolerance to detect zero singular values.

**Value**

An inversed matrix.

**See Also**

pinv from the pracma package

**Examples**

```
m <- cor(iris[1:4])
matrix_inverse(m)
```

---

simulate_simpson	<i>Simpson's paradox dataset simulation</i>
------------------	---

---

**Description**

Simpson's paradox, or the Yule-Simpson effect, is a phenomenon in probability and statistics, in which a trend appears in several different groups of data but disappears or reverses when these groups are combined.

**Usage**

```
simulate_simpson(n = 100, r = 0.5, groups = 3, difference = 1)
```

**Arguments**

n	The number of observations for each group to be generated.
r	A value or vector corresponding to the desired correlation coefficients.
groups	Number of groups.
difference	Difference between groups.

**Value**

A dataset.

**Examples**

```
data <- simulate_simpson(n = 100, groups = 5, r = 0.5)

library(ggplot2)
ggplot(data, aes(x = V1, y = V2)) +
  geom_point(aes(color = Group)) +
  geom_smooth(aes(color = Group), method = "lm") +
  geom_smooth(method = "lm")
```

---

z\_fisher

*Fisher z-transformation*

---

**Description**

The Fisher z-transformation converts the standard Pearson's  $r$  to a normally distributed variable  $z'$ . It is used to compute confidence intervals to correlations. The  $z'$  variable is different from the z-statistic.

**Usage**

```
z_fisher(r = NULL, z = NULL)
```

**Arguments**

r, z	The r or the $z'$ value to be converted.
------	--

**Value**

The transformed value.

**References**

Zar, J.H., (2014). Spearman Rank Correlation: Overview. Wiley StatsRef: Statistics Reference Online. doi:10.1002/9781118445112.stat05964



**Examples**

```
z_fisher(r = 0.7)
z_fisher(z = 0.867)
```

# Index

`cor_test`, 6  
`cor_to_ci`, 10  
`cor_to_cov`, 11  
`cor_to_p` (`cor_to_ci`), 10  
`cor_to_pcor`, 12  
`cor_to_spcor` (`cor_to_pcor`), 12  
`correlation`, 2

`distance_mahalanobis`, 13

`is.cor`, 14  
`isSquare`, 14

`matrix_inverse`, 15  
`model_parameters`, 3, 7

`pcor_to_cor` (`cor_to_pcor`), 12

`simulate_simpson`, 15  
`spcor_to_cor` (`cor_to_pcor`), 12

`z_fisher`, 16